

# Attachment C

A

To whom it may concern,

Cameron M (DOB 1/1/2012) has been diagnosed with a seizure disorder, and has been prescribed a medicine that can be given in case of prolonged seizure. If he has a seizure over 5 minutes long, he can be given 8 mg of intranasal Midazolam. Please call his parents if he has a seizure before giving the medicine.

Thank you,

Dr. [REDACTED]

B

Place child's photo here

Seizure Emergency Care Plan and Medication Orders for School Childcare Settings

**PARENT/GUARDIAN complete and sign the top portion of form.**

Child Name: Cameron [redacted]	Birth date: [redacted]
Parent/Guardian Contact:	Phone:
Emergency Contact:	Phone:
School:	Grade:

Triggers:  tiredness  flashing lights  illness  hunger  temperature  Other: \_\_\_\_\_

Seizure Aura (if any): \_\_\_\_\_

Seizure history:  Convulsive  Focal  Absence Date of last known seizure \_\_\_\_\_

Describe:

Anti-seizure Medication Taken at Home	Common side effects
Valproic acid	headache, fatigue

Other Seizure Treatments/Special Diet Therapy:

I give permission for school personnel to share this information, follow this plan, administer medication and care for my child and, if necessary, contact our health care provider. I assume full responsibility for providing the school with prescribed medication and devices. I approve this Seizure Emergency Care Plan for my child.

Parent Signature	Date	Nurse/CCHC Signature	Date
------------------	------	----------------------	------

HEALTH CARE PROVIDER to complete all items, SIGN and DATE completed form.

IF YOU SEE THIS:	DO THIS:
<input checked="" type="checkbox"/> <b>Convulsive Generalized Tonic Clonic:</b> You will see loss of consciousness. Stiffening of the body. Rhythmic jerking movements. Convulsive seizures may last 1-5 minutes. The child may have a warning (aura) before the seizure. Sleepiness and confusion may occur after the seizure.	<ol style="list-style-type: none"> <li>1. Time the seizure and record observations.</li> <li>2. Keep calm. Provide reassurance. Remove bystanders.</li> <li>3. Protect head, keep airway clear, turn on side.</li> <li>4. Do not restrain or place anything in mouth.</li> <li>5. Call 911 if student is injured or has difficulty breathing.</li> <li>6. Call parent.</li> <li>7. Stay with student until recovered from seizure.</li> <li>8. Administer rescue treatments as marked below.</li> </ol>
<input type="checkbox"/> <b>Focal:</b> These seizures may begin with an aura. They may be partly alert or unconscious. You may see lip smacking, chewing, eye blinking, or picking at clothes. These seizures usually last 1-2 minutes.	<ol style="list-style-type: none"> <li>1. Time the seizure and record observations.</li> <li>2. Gently guide child away from danger.</li> <li>3. Stay with student and reassure them until recovered from seizure.</li> <li>4. Do not treat staring that is stopped by a touch/nudge.</li> <li>5. Call parent.</li> <li>6. Administer rescue treatments as marked below.</li> </ol>
<input checked="" type="checkbox"/> <b>Absence/Myoclonic:</b> You will see quick jerk then changes in alertness. May see eye flutter or small twitching. Usually last less than 10 seconds.	

**RESCUE TREATMENTS:**

Vagus Nerve Stimulator: Child does not have a VNS.

**If seizure lasts longer than 5 minutes, administer:**

Diastat: <input type="checkbox"/> Diastat: N/A	<input checked="" type="checkbox"/> Midazolam 8 mg (1.6 mL) in the nose	<input type="checkbox"/> Clonazepam: N/A
--	---	--

Multistep seizure rescue plan – Please see attached letter for details.

**OR If cluster of 6 or more seizures in 60 minutes, administer:**

diastat: <input type="checkbox"/> Diastat: N/A	<input checked="" type="checkbox"/> Midazolam 8 mg (1.6 mL) in the nose	<input type="checkbox"/> Clonazepam: N/A
--	---	--

Multistep seizure rescue plan – Please see attached letter for details.

If emergency medication is administered:  call 911 if seizure does not stop within 5 min of giving medication

Other:

If no emergency medication is at school and the child is experiencing seizures:  
 Call family to bring medications to school or pick up child. Call EMS if seizure lasts more than 5 minutes.

Accommodations: Always take seizure action plan and emergency medication for school activities, sports and field trips. Close adult supervision when swimming or climbing.